

Treatment of Mandibular Angle Fractures

Jung-Ho Lee

Department of Plastic and Reconstructive Surgery, College of Medicine, The Catholic University of Korea, Seoul, Korea

No potential conflict of interest relevant to this article was reported.

The management of mandibular angle fractures is often challenging and results in the highest complication rate among fractures of the mandible. In addition, the optimal treatment modality for angle fractures remains controversial. Traditional treatment protocols for angle fractures have involved rigid fixation with intraoperative maxillomandibular fixation (MMF) to ensure absolute stability. However, more recently, non-compression miniplates have gained in popularity and the use of absolute intraoperative MMF as an adjunct to internal fixation has become controversial. In this article, the history of, and current trends in, the treatment of mandibular angle fractures will be briefly reviewed. In addition, issues regarding the management of the third molar tooth will be discussed.

Keywords: Mandibular fractures / Fracture fixation, internal / Molar, third

INTRODUCTION

Due to its prominence, the mandible is frequently involved in fractures of the facial bones. A retrospective review published in 2007 showed that in patients with a facial fracture, mandibular fractures ranked second (23.3%) to nasal bone fractures (58.6%) [1]. Among mandibular fractures, the incidence of angle fracture is relatively high (27%–30%) because the cross-sectional area is relatively thin within the angle, and also because of the presence of the third molar tooth [2]. Although many studies in the literature have suggested technical options to treat such fractures, the data are variable and the ideal treatment modality remains controversial. In addition, the reported complication rates range from 0–32% [3–10].

In this study, current trends in the treatment of isolated mandibular angle fractures will be briefly reviewed.

VARIOUS TECHNIQUES

Various techniques, including closed reduction, open reduction

by non-rigid fixation with wire, and open reduction and rigid internal fixation with plates or lag screws, have been reported for the management of mandibular angle fractures [11]. Although open reduction and internal fixation were first introduced in 1888, external techniques have predominated due to the poor treatment results associated with the corrosion and fatigue of metal plates.

In the 1960s, with the introduction of Viltallium compression plating by Luhr [12], internal fixation began to gain popularity. In the 1970s, the AO Foundation/Association for the Study of Internal Fixation (AO/ASIF) developed bone healing techniques that involved compression via dynamic compression plating. They stressed the need for absolute stability to prevent fragment mobility and to ensure primary bone healing [13]. Hence, for treatment of angle fractures, the original AO technique involved the placement of double plates along the superior and inferior borders of the mandible.

Simultaneously, Michelet et al. [14] began experimenting with monocortical non-compression plates for mandibular angle fractures. Champy et al. [15] showed that absolute rigid fixation was not mandatory for the healing of mandibular fractures and recommended the use of a single non-compression miniplate on the superior border of the mandible for angle fractures. With this technique, safe placement of the plate without causing damage to

Correspondence: Jung-Ho Lee

Department of Plastic Surgery, Bucheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, 327 Sosa-ro, Bucheon 14647, Korea
E-mail: tfm0822@catholic.ac.kr

Received January 2, 2017 / Revised April 13, 2017 / Accepted April 16, 2017

the roots of the teeth was possible because the size of the plate was small and the screws were monocortical. Subsequently, Ellis et al. [6,7,10] identified a trend towards increasing complication rates with increasing rigidity of fixation. This appeared to be because rigid fixation with a smaller number of plates required less periosteal stripping, thereby reducing disruption of the blood supply [16]. A recent systemic meta-analysis also showed that the single miniplate technique was statistically significantly superior compared with the two miniplate technique with regard to the incidence of postoperative complications [17].

However, many biomechanical tests continue to demonstrate the stability of two plates compared to single plate fixation for mandibular angle fractures and some investigators have advocated the use of two plate fixation for angle fractures [18-20]. In these studies, bony gaps were observed along the inferior border and such movement of the fracture was considered to contribute to subsequent complications, such as infection.

In spite of these conflicting results, the concepts underlying the management of mandibular angle fractures continue to evolve and a transition from large, dual compression plates to a single miniplate appears to be evident. A survey by the North American and European AO/ASIF regarding treatment modalities for mandibular angle fractures showed that about 51% of responders preferred the Champy technique while 22% preferred the placement of multiple plates. Interestingly, surgeons who treated more than 10 mandibular fractures per year favored the Champy technique over the tension band and bicortical plate combination, while surgeons who treated less than 10 mandibular fractures per year favored the tension band and bicortical compression plate combination over the Champy technique. Although superior placement of a single plate is generally preferred, an inferior border plate is indicated if adequate bone is lacking at the superior border (comminuted fracture), or if there is a history of previously failed hardware or a pathologic fracture is present [17].

Another controversy involves the use of postoperative maxillo-mandibular fixation (MMF). In a retrospective study of 287 patients, Valentino and Marentette [21] found that the addition of MMF did not significantly alter complication rates. Similarly, Kumar et al. [22] found no significant differences in outcomes or

complications between internal fixation with immediate release and internal fixation with 5–7 days of MMF. However, postoperative MMF still seems to have several advantages, including allowing the undisturbed healing of the intraoral incision, stabilizing the occlusion, and encouraging patients to become accustomed to a liquid diet [16].

The final point to consider is the management of the third molar tooth. Traditionally, the third molar tooth in the line of an angle fracture was known to be associated with an increased risk of infection, because intraoral communication via the periodontal ligament promotes the ingress of bacteria-laden saliva to the fracture site [16,23,24]. The absolute indications for the removal of the third molar tooth are as follows: (1) non-restorable damage to the tooth substance; (2) grade II or III mobility due to chronic periodontitis; (3) the presence of caries with periapical pathology; and (4) a displaced or extracted tooth preventing reduction [25].

On the other hand, some authors have advocated that the tooth in the line of the fracture should be preserved [25]. The preserved tooth can help with the repositioning of the fracture segments and can be used later on as an abutment for prosthesis placement. In addition, extraction might cause trauma and compounding of the fracture, which precludes rigid fixation. A recent systemic review and meta-analysis also showed that there was no significant statistical difference between removing or retaining the tooth in the line of the fracture with respect to the occurrence of postoperative infection [26].

These conflicting results indicate that the occurrence of postoperative infection does not depend solely on the status of the third molar tooth, but on many other factors (e.g., adequacy of the fixation, administration of adequate antibiotics, socioeconomic condition of the patient, oral hygiene, postoperative compliance of the patient, etc.).

CONCLUSION

In conclusion, although a trend does exist, the definitive treatment option for mandibular angle fractures remains elusive. This is perhaps to be expected because there are numerous confounding preoperative, intraoperative, and postoperative factors that

can affect surgical outcomes. It is therefore important to understand the pros and cons of each treatment option and individualize it according to the unique condition of the patient.

REFERENCES

- Allareddy V, Allareddy V, Nalliah RP. Epidemiology of facial fracture injuries. *J Oral Maxillofac Surg* 2011;69:2613-8.
- Haug RH, Prather J, Indresano AT. An epidemiologic survey of facial fractures and concomitant injuries. *J Oral Maxillofac Surg* 1990;48:926-32.
- Wagner WF, Neal DC, Alpert B. Morbidity associated with extraoral open reduction of mandibular fractures. *J Oral Surg* 1979;37:97-100.
- Chuong R, Donoff RB, Guralnick WC. A retrospective analysis of 327 mandibular fractures. *J Oral Maxillofac Surg* 1983;41:305-9.
- Iizuka T, Lindqvist C. Rigid internal fixation of fractures in the angular region of the mandible: an analysis of factors contributing to different complications. *Plast Reconstr Surg* 1993;91:265-71.
- Ellis E 3rd, Walker L. Treatment of mandibular angle fractures using two noncompression miniplates. *J Oral Maxillofac Surg* 1994;52:1032-6.
- Ellis E 3rd. Treatment methods for fractures of the mandibular angle. *Int J Oral Maxillofac Surg* 1999;28:243-52.
- James RB, Fredrickson C, Kent JN. Prospective study of mandibular fractures. *J Oral Surg* 1981;39:275-81.
- Schmelzeisen R, McIlff T, Rahn B. Further development of titanium miniplate fixation for mandibular fractures: experience gained and questions raised from a prospective clinical pilot study with 2.0 mm fixation plates. *J Craniomaxillofac Surg* 1992;20:251-6.
- Ellis E 3rd, Walker LR. Treatment of mandibular angle fractures using one noncompression miniplate. *J Oral Maxillofac Surg* 1996;54:864-71.
- Gear AJ, Apasova E, Schmitz JP, Schubert W. Treatment modalities for mandibular angle fractures. *J Oral Maxillofac Surg* 2005;63:655-63.
- Luhr HG. Vitallium Luhr systems for reconstructive surgery of the facial skeleton. *Otolaryngol Clin North Am* 1987;20:573-606.
- Schierle HP, Schmelzeisen R, Rahn B, Pytlik C. One- or two-plate fixation of mandibular angle fractures? *J Craniomaxillofac Surg* 1997;25:162-8.
- Michelet FX, Deymes J, Dessus B. Osteosynthesis with miniaturized screwed plates in maxillo-facial surgery. *J Maxillofac Surg* 1973;1:79-84.
- Champy M, Lodde JP, Schmitt R, Jaeger JH, Muster D. Mandibular osteosynthesis by miniature screwed plates via a buccal approach. *J Maxillofac Surg* 1978;6:14-21.
- Mehra P, Murad H. Internal fixation of mandibular angle fractures: a comparison of 2 techniques. *J Oral Maxillofac Surg* 2008;66:2254-60.
- Al-Moraissi EA, Ellis E 3rd. What method for management of unilateral mandibular angle fractures has the lowest rate of postoperative complications? A systematic review and meta-analysis. *J Oral Maxillofac Surg* 2014;72:2197-211.
- Choi BH, Kim KN, Kang HS. Clinical and in vitro evaluation of mandibular angle fracture fixation with the two-miniplate system. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995;79:692-5.
- Dichard A, Klotch DW. Testing biomechanical strength of repairs for the mandibular angle fracture. *Laryngoscope* 1994;104:201-8.
- Levy FE, Smith RW, Odland RM, Marentette LJ. Monocortical miniplate fixation of mandibular angle fractures. *Arch Otolaryngol Head Neck Surg* 1991;117:149-54.
- Valentino J, Marentette LJ. Supplemental maxillomandibular fixation with miniplate osteosynthesis. *Otolaryngol Head Neck Surg* 1995;112:215-20.
- Kumar I, Singh V, Bhagol A, Goel M, Gandhi S. Supplemental maxillomandibular fixation with miniplate osteosynthesis-required or not? *Oral Maxillofac Surg* 2011;15:27-30.
- Ellis E 3rd. Outcomes of patients with teeth in the line of mandibular angle fractures treated with stable internal fixation. *J Oral Maxillofac Surg* 2002;60:863-5.
- Tate GS, Ellis E 3rd, Throckmorton G. Bite forces in patients treated for mandibular angle fractures: implications for fixation recommendations. *J Oral Maxillofac Surg* 1994;52:734-6.
- Rai S, Pradhan R. Tooth in the line of fracture: its prognosis and its effects on healing. *Indian J Dent Res* 2011;22:495-6.
- Bobrowski AN, Sonogo CL, Chagas Junior OL. Postoperative infection associated with mandibular angle fracture treatment in the presence of teeth on the fracture line: a systematic review and meta-analysis. *Int J Oral Maxillofac Surg* 2013;42:1041-8.